

GLACIERS

Global glacier change in the 21st century: Every increase in temperature matters

David R. Rounce^{1,2*}, Regine Hock^{2,3}, Fabien Maussion⁴, Romain Hugonnet^{5,6,7}, William Kochtitzky^{8,9}, Matthias Huss^{5,6,10}, Etienne Berthier⁷, Douglas Brinkerhoff¹¹, Loris Compagno^{5,6}, Luke Copland⁸, Daniel Farinotti^{5,6}, Brian Menounos^{12,13}, Robert W. McNabb¹⁴

Glacier mass loss affects sea level rise, water resources, and natural hazards. We present global glacier projections, excluding the ice sheets, for shared socioeconomic pathways calibrated with data for each glacier. Glaciers are projected to lose $26 \pm 6\%$ ($+1.5^\circ\text{C}$) to $41 \pm 11\%$ ($+4^\circ\text{C}$) of their mass by 2100, relative to 2015, for global temperature change scenarios. This corresponds to 90 ± 26 to 154 ± 44 millimeters sea level equivalent and will cause 49 ± 9 to $83 \pm 7\%$ of glaciers to disappear. Mass loss is linearly related to temperature increase and thus reductions in temperature increase reduce mass loss. Based on climate pledges from the Conference of the Parties (COP26), global mean temperature is projected to increase by $+2.7^\circ\text{C}$, which would lead to a sea level contribution of 115 ± 40 millimeters and cause widespread deglaciation in most mid-latitude regions by 2100.

Glaciers, here referring to all glacial land ice excluding the Greenland and Antarctic ice sheets, are responsible for $21 \pm 3\%$ of sea level rise from 2000 to 2019, contributing 0.74 ± 0.04 mm sea level equivalent (SLE) yr^{-1} (1). Projections suggest this contribution could increase to 2.5 mm SLE yr^{-1} by 2100 (2). Glaciers are also a critical water resource for ~ 1.9 billion people (3), and projected losses will alter water availability impacting annual and seasonal runoff (4). Glacier-related hazards, including glacier outburst floods, are also expected to change in frequency and magnitude over the next century as a result of mass loss (5). Projecting the magnitude, spatial pattern, and timing of glacier mass loss is therefore essential to support climate adaptation and mitigation efforts for communities ranging from the coast to the high mountains.

Previous projections of glacier mass loss from the glacier model intercomparison project (GlacierMIP) (2) estimated glacier contribution to sea level rise for ensembles of representative concentration pathways (RCPs), and

results were extended to shared socioeconomic pathways (SSPs) using statistical models of these simulations (6). GlacierMIP provided these projections at regional scales based on simulations from 11 glacier evolution models that varied with respect to the complexity of model physics, simulated physical processes, model calibration, spatial resolution, and modeling domain. Calibration data varied from in situ measurements of less than 300 of the world's more than 215,000 glaciers to regional geodetic and/or gravimetric mass balance observations. Furthermore, only one global model simulated glacier dynamics using a flowline model (7), whereas all others relied on empirical volume-area scaling or parameterizations of mass redistribution; only one model accounted for frontal ablation (i.e., the sum of iceberg calving and submarine melt) of marine-terminating glaciers (8), whereas all others treated any glacier as land-terminating; further, no global model accounted for debris cover. Existing multimodel projections (2, 6, 9) are thus limited to regional scales and neglect key physical processes controlling glacier mass loss.

We produce a set of global glacier projections for every glacier on Earth for SSPs from 2015 to 2100 by leveraging global glacier mass balance data (1) and near-global frontal ablation data (10–13). To provide policy-relevant scenarios, our projections are grouped based on mean global temperature increases by the end of the 21st century compared with pre-industrial levels to explicitly link differences in glacier mass loss, sea level rise, and the number of glaciers that vanish in response to changes in mean global temperature. Our glacier evolution model, a hybrid of the Python Glacier Evolution Model (PyGEM) (14, 15) and Open Global Glacier Model (OGGM) (7), enables us to produce global glacier projections that explicitly account for glacier dynamics using a

flowline model (7) based on the shallow-ice approximation (16), the effects of debris thickness on sub-debris melt rates (17), and frontal ablation (8). Our estimates of glacier contribution to sea level rise also account for the $\sim 15\%$ of ice from marine-terminating glaciers that is already below sea level (18). Projections are also reported for SSPs and RCPs to highlight differences compared with previous studies.

Projections of policy-relevant scenarios

The Paris Agreement, adopted in 2015 by 195 countries, agreed to keep the increase in global mean temperature by the end of the 21st century relative to preindustrial levels below 2°C , and that efforts should be made to limit the temperature change to 1.5°C . This target was kept alive in the Glasgow Agreement adopted by the Conference of the Parties (COP26) in 2021. To evaluate the sensitivity of glaciers to global mean temperature increases, the glacier projections are aggregated into $+1.5^\circ\text{C}$, $+2^\circ\text{C}$, $+3^\circ\text{C}$, and $+4^\circ\text{C}$ temperature change scenarios by 2100 relative to preindustrial levels (Fig. 1).

Globally, glaciers are projected to lose $26 \pm 6\%$ ($+1.5^\circ\text{C}$) to $41 \pm 11\%$ ($+4^\circ\text{C}$) of their mass by 2100, relative to 2015 [ensemble median $\pm 95\%$ confidence interval (CI)]. This mass loss would increase mean sea level by 90 ± 26 mm SLE under the $+1.5^\circ\text{C}$ scenario and 99 ± 31 mm SLE under the $+2^\circ\text{C}$ scenario. The higher temperature change scenarios of $+3^\circ\text{C}$ and $+4^\circ\text{C}$ lead to contributions of 125 ± 39 and 154 ± 44 mm SLE, respectively, highlighting a 71% increase between the $+1.5^\circ\text{C}$ and $+4^\circ\text{C}$ scenarios.

The rate of sea level rise from glacier mass loss near the end of the 21st century ranges from 0.70 ± 0.45 to 2.23 ± 1.08 mm SLE yr^{-1} depending on the temperature change scenario (fig. S1). For $+1.5^\circ\text{C}$, the rate of sea level rise peaks at 1.29 ± 0.59 mm SLE yr^{-1} around 2035 and declines thereafter whereas the rate for $+4^\circ\text{C}$ steadily increases for the remainder of this century. Similar trends are observed in the area-averaged mass loss rate, where the maximum loss rate of 0.82 ± 0.36 m water equivalent (w.e.) yr^{-1} occurs around 2035 before diminishing to 0.59 ± 0.34 m w.e. yr^{-1} at the end of the century for the $+1.5^\circ\text{C}$ scenario; the mass loss rate continuously increases to 2.02 ± 1.30 m w.e. yr^{-1} by the end of the century for the $+4^\circ\text{C}$ scenario (Fig. 1I). Even if the global mean temperature change is limited to $+1.5^\circ\text{C}$, we estimate that 104,000 $\pm 20,000$ glaciers ($49 \pm 9\%$ of the total inventoried) will disappear by 2100 and at least half of those will be lost before 2050 (Fig. 1E). Most of the glaciers projected to disappear are <1 km² (Fig. 2) but regardless of their small size, their disappearance may still negatively affect local hydrology, tourism, glacier hazards, and cultural values (19). Glaciers projected to disappear represent 2 to 8% of the glacier contribution to sea level rise depending on the temperature change scenario.

¹Department of Civil and Environmental Engineering, Carnegie Mellon University, Pittsburgh, PA, USA. ²Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK, USA. ³Department of Geosciences, University of Oslo, Oslo, Norway. ⁴Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria. ⁵Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zürich, Zurich, Switzerland. ⁶Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland. ⁷LEGOS, Université de Toulouse, CNES, CNRS, IRD, UPS, Toulouse, France. ⁸Department of Geography, Environment and Geomatics, University of Ottawa, Ottawa, Ontario, Canada. ⁹School of Marine and Environmental Programs, University of New England, Biddeford, ME, USA. ¹⁰Department of Geosciences, University of Fribourg, Fribourg, Switzerland. ¹¹Department of Computer Science, University of Montana, Missoula, MT, USA. ¹²Geography Earth and Environmental Sciences, University of Northern British Columbia, Prince George, BC, Canada. ¹³Hakai Institute, Campbell River, BC, Canada. ¹⁴School of Geography and Environmental Sciences, Ulster University, Coleraine, UK.

*Corresponding author. Email: drounce@cmu.edu

Regional mass changes

Regional variations exist in the glacier mass change projections (Fig. 3). Alaska is the largest regional contributor to global mean sea level rise from 2015 to 2100 (fig. S2), peaking at 0.33 to 0.44 mm SLE yr^{-1} between 2030 and 2060 depending on the temperature change scenario, before decreasing to 0.13 to 0.28 mm SLE yr^{-1} by 2100 (fig. S1). Greenland Periphery, Antarctic and Subantarctic, Arctic Canada North, and Arctic Canada South contribute 12, 10, 10, and 9% to projected sea level rise, respectively. Collectively, these five regions account for 60 to 65% of the total glacier contribution to sea level rise. For Greenland Periphery, Arctic Canada North, and Arctic Canada South, the rate of the contribution to sea level rise is almost insensitive to temperature change below +2°C but steadily increases through 2100 for the other temperature change scenarios. For the +3°C and +4°C scenarios, the rate of sea-level rise from Greenland Periphery, Antarctic and Subantarctic, and Arctic Canada North each nearly equal or exceed Alaska near the end of the century, with Antarctic and Subantarctic and Arctic Canada North accelerating throughout the 21st century. Because projected glacier mass loss includes both the instantaneous response of glaciers to climate forcing and the delayed response based on the extent of disequilibrium to longer-term climatic conditions (20), these regions with large glaciers will continue losing mass beyond 2100, especially for higher temperature change scenarios.

Western Canada and US, South Asia East, Scandinavia, North Asia, Central Europe, Low Latitudes, Caucasus and Middle East, and New Zealand are projected to lose 60 to 100% of their glacier mass depending on the temperature change scenario (Fig. 3 and fig. S3). The temperature change scenario thus has a major impact on the mass loss, in some cases determining whether the complete deglaciation of regions occurs by the end of the 21st century. Although these regions are not significant contributors to sea level rise, people in these regions will need to adapt to changes in seasonal and annual runoff as the additional water provided by glacier net mass loss will decline before 2050 as the glaciers retreat (figs. S5 and S8). In High Mountain Asia, the timing of maximum rates of mass loss varies, with South Asia East peaking between 2025 and 2030, Central Asia between 2035 and 2055, and South Asia West between 2050 and 2075, depending on the temperature change scenario.

Regional sensitivity to temperature change

The sensitivity of the glacierized regions to changes in global mean temperature depends on the region's current glacier mass and mass change rates; regional temperature anomalies relative to the global mean (Fig. 4), such




Fig. 1. Projected global glacier changes for scenarios of global mean temperature change. (A and B) Mass remaining, (C and D) area remaining, (E and F) glaciers remaining, (G and H) sea level rise (SLR) contributed from glaciers, and (I and J) area-averaged mass change rate for all glaciers globally. Projections are shown from 2015 to 2100 (left panels), and at 2100 (right panels). Values in [(A) to (H)] are relative to 2015. Colors depict the global mean temperature change scenarios (left panels) and the SSPs corresponding to the global temperature changes (right panels). The number (n) of glacier projections with different general circulation models (GCMs) and SSPs that fall into each temperature change scenario is shown in the legend. Lines (left panels) show the ensemble median and shading indicates the 95% CI for each temperature change scenario.




Fig. 2. Percent of glaciers projected to vanish between 2015 and 2100 for global temperature change scenarios sorted by size. The glaciers are binned according to their initial glacier area and the numbers below each bin (shown in gray) refer to the percentage of the total number of glaciers in 2015 in each bin.

as those associated with Arctic amplification (21); the climatic setting (maritime versus continental); sensitivity to precipitation falling as rain instead of snow; and elevation feedbacks due to different types of glaciers (e.g., ice caps versus valley glaciers) (22). Projected mass

loss is linearly related to global mean temperature increase, especially for larger glacierized regions, consistent with a recent study (6). This strong relationship highlights that every fraction of a degree of temperature increase substantially affects glacier mass loss. The




Fig. 3. Regional glacier mass change and contributions to sea level rise from 2015 to 2100. Discs show global and regional projections of glacier mass remaining by 2100 relative to 2015 for global mean temperature change scenarios. Discs are scaled based on each region's contribution to global mean sea level rise from 2015 to 2100 for the +2°C scenario by 2100 relative to preindustrial levels, and nested rings are colored by temperature change scenarios showing normalized mass remaining in 2100. Regional sea level rise contributions >1 mm SLE for the +2°C scenario are printed in the center of each disc. The horizontal bars below each disc show time series of area-averaged annual mass balance from 2015 to 2100 for +1.5°C (top bar) and +3°C (bottom bar) scenarios. The colorbar is saturated at -2.5 m w.e., but minimum annual values reach -4.2 m w.e. in Scandinavia. Time series of regional relative mass change and regional area-averaged mass change are shown in figs. S3 and S4.

smallest glacierized regions by mass, including Central Europe, Scandinavia, Caucasus and Middle East, North Asia, Western Canada and US, Low Latitudes, and New Zealand, will experience near-complete deglaciation around +3°C. These regions are thus highly sensitive to global mean temperature increases between 1.5 and 3°C and have a nonlinear response above 3°C of warming.

The strength of the linear relationship varies among regions, which reflects differences in the regional temperature anomalies from the ensemble of GCMs (evident from the larger standard deviations given in Fig. 4 and fig. S9). Regions like Alaska, Southern Andes, and Central Asia have less scatter, indicating less variation in the regional temperature anomaly and thereby a more consistent response to climate forcing (mean $R^2 = 0.78$). Other regions like the Russian Arctic, Svalbard, and Iceland have more variation in the regional temperature anomaly and thus a weaker linear relationship (mean $R^2 = 0.50$) as well as considerable variation in projected precipitation (fig. S10). Future work using regional climate projections may better resolve high-mountain cli-

matic conditions and refine projections in these regions.

Spatially resolved projections at the glacier scale

Our projections reveal notable spatial variation in glacier mass loss at the local scale for the temperature change scenarios (Fig. 5). All regions are projected to lose some glaciers completely, primarily smaller ice masses, with the higher temperature change scenarios revealing significantly more mass loss and the deglaciation of greater areas (figs. S11 to S13). Although Central Europe, Caucasus and Middle East, North Asia, and Western Canada and US are projected to experience widespread deglaciation for the +2°C scenario, our results also reveal where remaining glaciers will be concentrated at the end of this century. Besides the Karakoram and Kunlun in High Mountain Asia, the remaining mass is primarily located in southeastern Alaska, Arctic Canada North, Svalbard, the Russian Arctic, Greenland Periphery, and Antarctic and Subantarctic. Given that these regions constitute a significant number of marine-terminating glaciers,

accounting for frontal ablation is critical over the next century and beyond.

Importance of marine-terminating glaciers

Marine-terminating glaciers represent 40% of the total present-day global glacier area (23), and this percentage reaches 99% for the Antarctic and Subantarctic region. Most previous global glacier projections do not explicitly account for frontal ablation (2) and instead implicitly account for it by increasing melt rates, thereby poorly accounting for dynamical feedbacks associated with the glacier's evolution. Our model couples a frontal ablation parameterization with a flowline model and uses a state-of-the-art calibration scheme, ice thickness inversion method, and geodetic mass balance and frontal ablation calibration data (see Methods). These features enable us to project changes of individual marine-terminating glaciers and determine if and when they become land-terminating (fig. S14). Separate simulations including and excluding frontal ablation, with model parameters calibrated separately for both, are used to quantify the impact of accounting for frontal ablation on projections.

Counterintuitively, we estimate that accounting for frontal ablation reduces the glacier contribution to mean sea level rise from 2015 to 2100 by 2% for each temperature change scenario, compared with models not including frontal ablation. From 2015 to 2100 frontal ablation accounts for $91 \pm 10 \text{ Gt yr}^{-1}$ (+1.5°C) to $88 \pm 8 \text{ Gt yr}^{-1}$ (+4°C) of the total glacier mass loss globally (figs. S15 to S18). For the +2°C scenario, the rate of mass loss due to frontal ablation diminishes over the century from $115 \pm 11 \text{ Gt yr}^{-1}$ in 2000 to 2020 to $75 \pm 8 \text{ Gt yr}^{-1}$ in 2080 to 2100. Diminished mass losses from frontal ablation of marine-terminating glaciers reflect their thinning, retreat onto land (44 to 57% of all marine-terminating glaciers) (fig. S19), and reduced ice flux into the ocean, which occurs for all temperature change scenarios. The relative contribution of frontal ablation to total ablation (i.e., frontal ablation plus melt) ranges from 11% (+1.5°C) to 8% (+4°C) for 2015 to 2100, diminishing for higher temperature change scenarios due to increases in melt. Regionally, the relative contribution of frontal ablation for all temperature change scenarios is greatest in Antarctic and Subantarctic (34%), the Russian Arctic (34%), and Svalbard (17%) (figs. S15 to S18).

The impact of not accounting for frontal ablation on relative mass loss (i.e., glacier mass loss by 2100 relative to 2015) varies greatly by region (fig. S20). For Alaska and Svalbard, excluding frontal ablation increases relative mass loss at 2100 by 2 to 8% depending on the temperature change scenario. The Russian Arctic varies from a 2% reduction (+1.5°C) to a 5% increase (+4°C). Arctic Canada, Greenland Periphery, and Southern Andes see almost

no difference ($\pm 2\%$), and Antarctic and Subantarctic see a 0 to 2% decrease in relative mass loss. These results highlight the complex response of marine-terminating glaciers, which are dependent on the frontal ablation rate, glacier geometry, and surface mass balance. In the Antarctic and Subantarctic, we find excluding frontal ablation decreases the regional relative mass loss, as mass loss due to frontal ablation is greater than the increased melt when frontal ablation is excluded. Conversely, in Alaska and Svalbard, the regional relative mass loss increases when frontal ablation is excluded, as mass loss due to frontal ablation is less than the increased melt when frontal ablation is excluded.

Importance of debris-covered glaciers

Debris currently covers 4 to 7% of the global glacier area (24, 25). A thin layer of debris (<3 to 5 cm) enhances surface melt, whereas a thick layer insulates the underlying ice and reduces melt (26). The spatial distribution of debris thickness can cause debris-covered glaciers to develop stagnant glacier tongues and eventually separate from the active part of the glacier (27, 28). Our representation of debris and glacier dynamics enables us to simulate these complex feedbacks, including reduced melt at glacier termini where debris is thick (fig. S21). We thus produce a set of global glacier projections that account for debris and compare these to separate simulations that exclude debris (i.e., treating the debris as clean ice) to quantify the insulating effect that debris has on glacier projections.

The impact of debris on relative mass loss varies greatly spatially and temporally (fig. S22) with the most significant differences occurring around mid-century in New Zealand and South Asia East. In these regions, the insulating effect of debris reduces net mass loss by 9 to 13% depending on the temperature change scenario, although the differences are less than 5% by 2100. Alaska, the largest region by mass with considerable debris cover (>5% by area), sees a reduction of 5% around 2060 and 3% by 2100. Other regions with considerable debris cover (>5% by area), including Western Canada and US, Central Europe, Caucasus and Middle East, and Low Latitudes, see a reduction in mass loss of less than 5% around mid-century and no difference ($\pm 1\%$) by 2100. The inclusion of debris thus delays mass loss over the century especially at local scales but has little impact on sea level rise and the number of glaciers lost by 2100. The limited impact in most regions shows that the insulating effect of debris is unable to offset the increased melt for the various temperature change scenarios.

Comparison with previous projections

For comparison with recent multimodel studies (2, 6), we also report our projections for the

RCPs and SSPs. Our global projections of glacier contribution to sea level rise for 2015 to 2100 range from 90 ± 36 mm SLE (RCP2.6) to 163 ± 53 mm SLE (RCP8.5) and 98 ± 38 mm SLE (SSP1-2.6) to 166 ± 83 mm SLE (SSP5-8.5), respectively (Table 1). These projections include a correction (reduction) of 17 to 24 mm SLE, which accounts for the mass loss of ice from marine-terminating glaciers that is below sea level and therefore will not contribute to global mean sea level rise—an important difference compared with current multimodel studies (2, 6), which do not account for this.

Even with this correction, for the low emissions scenarios our RCP2.6 projections are 11 mm SLE (14%) greater than that of Marzeion *et al.* (2), and our SSP1-2.6 projections are 18 mm SLE (23%) greater than that of Edwards *et al.* (6). For the mid-range (RCP4.5 and SSP2-4.5) and high (RCP8.5, SSP5-8.5) emissions scenarios, our projections are within ± 7 mm SLE of both studies.

Not correcting for the loss of ice below sea level, our projections of glacier contribution to sea level rise from 2015 to 2100 are 11 to 44% greater than these multimodel estimates




Fig. 4. Fraction of global and regional mass remaining at 2100, relative to 2015, as a function of global mean temperature change by 2100 relative to preindustrial levels. Each marker represents results from one GCM and SSP. Numbers indicate median temperature anomalies (\pm standard deviation) ($^{\circ}$ C) over glacierized areas, relative to the mean temperature change over the entire globe at 2100 relative to preindustrial levels, for all GCMs and scenarios, and the glacier mass at 2015 (10^3 Gt). Negative values indicate that some regions warm less than the global average. Regions are ordered by their total mass loss.




Fig. 5. Spatial distribution of glacier mass remaining by 2100 for the +2°C scenario. The ensemble median glacier mass remaining by 2100 (relative to 2015) for the +2°C (above preindustrial levels) global mean temperature change scenario. Tiles are aggregated by 1° by 1° below 60° latitude, 2° by 1° between 60° and 74° latitude and 2° by 2° above 74° latitude to represent ~10,000 km² each. Circles are scaled based on simulated glacierized area in

2015 and are colored by normalized mass remaining. Regions that have experienced complete deglaciation by 2100 are shown in white and outlined in black. High Mountain Asia refers to Central Asia, South Asia West, and South Asia East. Specific subregions are noted by labels on the bottom of inset figures. Additional temperature change scenarios (+1.5°C, +3°C, and +4°C) are shown in figs. S11 to S13.

Downloaded from https://www.science.org at University of Northern British Columbia on January 05, 2023

Table 1. Projected global glacier mass loss and glacier contribution to sea level rise. Results are shown for RCP and SSP scenarios at 2100, relative to 2015, from this study and recent multimodel studies (2, 6). “Uncorrected” refers to projections that assume mass losses below sea level contribute to sea level rise, consistent with assumptions in recent multimodel studies. Note that uncertainty associated with the multimodel studies is expressed as 90% CI, whereas this study reports ensemble median and 95% CI. Regional comparisons are shown in tables S1 and S2.

Global glacier contribution to sea level rise from 2015 to 2100 (mm SLE)

Study	RCP2.6	RCP4.5	RCP8.5	SSP1-2.6	SSP2-4.5	SSP5-8.5
This study	90 ± 36	114 ± 44	163 ± 53	98 ± 38	116 ± 51	166 ± 83
This study (uncorrected)	106 ± 37	132 ± 47	187 ± 61	115 ± 42	135 ± 57	192 ± 97
Marzeion <i>et al.</i> (2)	79 ± 57	119 ± 66	159 ± 86	-	-	-
Edwards <i>et al.</i> (6)	-	-	-	80 ± 35	119 ± 39	159 ± 47

Global glacier mass loss, relative to 2015 (%)

Study	RCP2.6	RCP4.5	RCP8.5	SSP1-2.6	SSP2-4.5	SSP5-8.5
This study	26 ± 8	31 ± 10	43 ± 13	28 ± 9	32 ± 12	44 ± 20
Marzeion <i>et al.</i> (2)	18 ± 13	27 ± 15	36 ± 20	-	-	-

(2, 6) for all emission scenarios. We attribute these differences to the global mass balance data we used for calibration, which include an accelerated trend in mass loss from 2000 to 2020 (1), as well as the improved representation of physical processes in our model.

Globally, we predict glaciers will lose 26 ± 8% (RCP2.6) to 43 ± 13% (RCP8.5) and 28 ± 9% (SSP1-2.6) to 44 ± 20% (SSP5-8.5) of their mass by 2100, relative to 2015. Our projected

relative mass losses are 4 to 8% greater than current multimodel estimates (2). Regionally, the most significant differences occur in Alaska, Arctic Canada South, South Asia East, and Southern Andes, where we predict 11 to 23% more relative mass loss (table S1). In Alaska, we estimate 22% (RCP2.6) to 23% (RCP8.5) more relative mass loss compared with the multimodel estimates (2) and find a peak in the net mass loss rate in the middle

of the century, in contrast to the peak net mass loss rate at the end of the century from the multimodel estimates (2).

A comparison of our projections from the ensembles of RCPs and SSPs used in this study reveals that glacier contribution to sea level rise is 2 to 9% greater for SSPs than the corresponding RCPs. These differences are a result of the SSPs simulating greater temperature increases for the same radiative forcing

as the RCPs (29, 30). Our ensembles reflect this higher warming sensitivity as the SSPs are on average 0.14 to 0.25°C warmer than their corresponding RCPs. Considering the high sensitivity of global and regional glacier mass loss to small temperature increases revealed by our study, the higher warming sensitivity of the SSPs will substantially affect the projected glacier contribution to sea level rise as well as the number of glaciers anticipated to be lost.

Summary and way forward

Our projections reveal a strong linear relationship between global mean temperature increase and glacier mass loss, with the smallest glacierized regions having a nonlinear relationship beyond +3°C as they experience near-complete deglaciation. This strong relationship at global and regional scales highlights that every increase in temperature has significant consequences with respect to glacier contribution to sea level rise, the loss of glaciers around the world, and changes to hydrology, ecology, and natural hazards. Regardless of the temperature change scenario, all regions will experience considerable deglaciation at local scales with roughly half of the world's glaciers by number projected to be lost by 2100, even if temperature increase is limited to +1.5°C. Based on the most recent climate pledges from COP26, global mean temperature is estimated to increase by +2.7°C (31), which would result in much greater glacier contribution to sea level rise (115 ± 40 mm SLE) and the near-complete deglaciation of entire regions including Central Europe, Western Canada and US, and New Zealand (Fig. 5 and figs. S11 to S13) compared with the Paris Agreement. The rapidly increasing glacier mass losses as global temperature increases beyond +1.5°C stresses the urgency of estab-

lishing more ambitious climate pledges to preserve the glaciers in these mountainous regions.

REFERENCES AND NOTES

- R. Hugonnet et al., *Nature* **592**, 726–731 (2021).
- B. Marzeion et al., *Earths Futur.* **8**, (2020).
- W. W. Immerzeel et al., *Nature* **577**, 364–369 (2020).
- M. Huss, R. Hock, *Nat. Clim. Chang.* **8**, 135–140 (2018).
- S. Harrison et al., *Cryosphere* **12**, 1195–1209 (2018).
- T. L. Edwards et al., *Nature* **593**, 74–82 (2021).
- F. Maussion et al., *Geosci. Model Dev.* **12**, 909–931 (2019).
- M. Huss, R. Hock, *Front. Earth Sci.* **3**, (2015).
- R. Hock et al., *J. Glaciol.* **65**, 453–467 (2019).
- B. Osmanoglu, M. Braun, R. Hock, F. J. Navarro, *Ann. Glaciol.* **54**, 111–119 (2013).
- B. Osmanoglu, F. J. Navarro, R. Hock, M. Braun, M. I. Corcuera, *Cryosphere* **8**, 1807–1823 (2014).
- M. Minowa, M. Schaefer, S. Sugiyama, D. Sakakibara, P. Skvarca, *Earth Planet. Sci. Lett.* **561**, 116811 (2021).
- W. Kochtitzky et al., *Nat. Commun.* **13**, 5835 (2022).
- D. R. Rounce et al., *J. Glaciol.* **66**, 175–187 (2020).
- D. R. Rounce, R. Hock, D. E. Shean, *Front. Earth Sci.* **7**, 331 (2020).
- K. Hutter, *J. Glaciol.* **27**, 39–56 (1981).
- D. R. Rounce et al., *Geophys. Res. Lett.* **48**, GL091311 (2021).
- D. Farinotti et al., *Nat. Geosci.* **12**, 168–173 (2019).
- R. Hock et al., in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, H.-O. Pörtner et al., Eds. (Cambridge University Press, 2019).
- H. Zekollari, M. Huss, D. Farinotti, *Cryosphere* **13**, 1125–1146 (2019).
- F. Pithan, T. Mauritsen, *Nat. Geosci.* **7**, 181–184 (2014).
- J. Bollbar, A. Rabatel, I. Gouttevin, H. Zekollari, C. Galiez, *Nat. Commun.* **13**, 409 (2022).
- RGI Consortium, Randolph glacier inventory - A dataset of global glacier outlines, Version 6.0, GLIMS (2017); <https://doi.org/10.7265/4m1f-gd79>
- D. Scherler, H. Wulf, N. Gorelick, *Geophys. Res. Lett.* **45**, 11798–11805 (2018).
- S. Herreid, F. Pellicciotti, *Nat. Geosci.* **13**, 621–627 (2020).
- G. Østrem, *Geogr. Ann.* **41**, 228–230 (1959).
- D. I. Benn et al., *Earth Sci. Rev.* **114**, 156–174 (2012).
- A.V. Rowan et al., *J. Geophys. Res. Earth Surf.* **126**, (2021).
- K. B. Tokarska et al., *Sci. Adv.* **6**, aaz9549 (2020).
- K. Wyser, E. Kjellström, T. Koenig, H. Martins, R. Döscher, *Environ. Res. Lett.* **15**, 054020 (2020).
- UNEP, “Emissions Gap Report 2021” (UNEP, 2021); <https://www.unep.org/emissions-gap-report-2021>.

ACKNOWLEDGMENTS

This work was supported in part by the high-performance computing and data storage resources operated by the Research Computing Systems Group at the University of Alaska Fairbanks Geophysical Institute. This text reflects only the author's view and funding agencies are not responsible for any use that may be made of the information it contains. **Funding:** This work was funded by the following: National Aeronautics and Space Administration grant 80NSSC20K1296 (to D.R. and Re.Ho.); National Aeronautics and Space Administration grant 80NSSC20K1595 (to D.R. and Re.Ho.); National Aeronautics and Space Administration grant 80NSSC17K0566 (to D.R. and Re.Ho.); National Aeronautics and Space Administration grant NNX17AB27G (to D.R. and Re.Ho.); Norwegian Research Council project #324131 (to Re.Ho.); Tula Foundation and Canada Research Chairs (to B.M.); National Sciences and Engineering Research Council of Canada (to B.M. and Lu.Co.); Vanier Graduate Scholarship (to W.K.); Swiss National Science Foundation project 184634 (to Ro.Hu., M.H., Lo.Co., and D.F.); ArcticNet Network of Centres of Excellence Canada (to Lu.Co.); University of Ottawa, University Research Chair program (to Lu.Co.); European Union's Horizon 2020 research and innovation programme grant 101003687 (to F.M.); Austrian Science Fund (FWF) grant P30256 (to F.M.); French Space Agency CNES (to E.B. and Ro.Hu.)

Author contributions: Conceptualization: D.R. and Re.Ho. Data curation: D.R. Formal analysis: D.R. Funding acquisition: D.R., Re.Ho., M.H., D.F., E.B., B.M., and Lu.Co. Investigation: D.R. Methodology: D.R., F.M., and Re.Ho. Project administration: D.R. and Re.Ho. Resources: D.R., F.M. (glacier data); Ro.Hu., M.H., E.B., D.F., B.M., and R.M. (mass balance data); Lo.Co. (climate data); W.K. and Lu.Co. (frontal ablation data); Software: D.R. (PyGEM); F.M. (OGGM); D.B. (emulators); Visualization: D.R., Re.Ho., and Ro.Hu. Writing – original draft: D.R. Writing – review and editing: all authors, especially Re.Ho. **Competing interests:** Authors declare that they have no competing interests. **Data and materials availability:** The datasets generated for this study can be found in the National Snow and Ice Data Center (NSIDC) at https://nsidc.org/data/hma2_ggp/versions/1. The model code is publicly available at <https://github.com/drounce/PyGEM> and <https://github.com/OGGM/oggm>. **License information:** Copyright © 2023 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.sciencemag.org/about/science-licenses-journal-article-reuse>

SUPPLEMENTARY MATERIALS

science.org/doi/10.1126/science.abo1324

Methods
References (32–41)
Figs. S1 to S28
Tables S1 to S5

Submitted 15 January 2022; accepted 14 November 2022
10.1126/science.abo1324

Global glacier change in the 21st century: Every increase in temperature matters

David R. RounceRegine HockFabien MaussionRomain HugonnetWilliam KochtitzkyMatthias HussEtienne BerthierDouglas BrinkerhoffLoris CompagnoLuke CoplandDaniel FarinottiBrian MenounosRobert W. McNabb

Science, 379 (6627), • DOI: 10.1126/science.abo1324

Melting away

Mountain glaciers, perennial ice masses excluding the Greenland and Antarctic ice sheets, are a critical water resource for nearly two billion people and are threatened by global warming. Rounce *et al.* projected how those glaciers will be affected under global temperature increases of 1.5° to 4°C, finding losses of one quarter to nearly one half of their mass by 2100 (see the Perspective by Aðalgeirsdóttir and James). Their calculations suggest that glaciers will lose substantially more mass and contribute more to sea level rise than current estimates indicate. —HJS

View the article online

<https://www.science.org/doi/10.1126/science.abo1324>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

Science (ISSN) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works